

Short Communication

Illegal capture and black market trade of sea turtles in Pisco, Peru: the never-ending story

Javier Quiñones¹, Sixto Quispe¹ & Oscar Galindo¹

¹Laboratorio Costero de Pisco, Instituto del Mar del Perú, Callao, Perú
Corresponding author: Javier Quiñones (javierantonioquinones@gmail.com)

ABSTRACT. The Pisco-San Andrés area (13°44'S, 76°13'W) in central Peru is known for a traditional historic sea turtle fishery. To determine if illegal captures and black market trade exist, we carried out bi-weekly sampling in dumpsites and coastal areas from 2009 to 2015. A total of 953 carapaces were encountered, which included mainly black turtles (*Chelonia mydas*, 92.2%) and to a lesser extent, olive ridley turtles (*Lepidochelys olivacea*, 4.3%), leatherback turtles (*Dermochelys coriacea*, 1.4%), and a single hawksbill turtle (*Eretmochelys imbricate*, 0.1%). The mean curved carapace length (CCL) was 59.1 for black turtles, 60.4 for olive ridleys and 113 cm for leatherbacks. For all species, most of turtles reported were juveniles and came largely from illegal captures (89%) and not from stranding reports (1.4%). Mean mortality was 8.1 carcasses km⁻¹ year⁻¹ at beaches and 160.2 carcasses year⁻¹ at dumpsites. Main consumed prey items in black turtles were silverside fish eggs (47.9%), *Chondracanthus* seaweed (31.4%) and *Paranthus* sp., anemone (16.2%). Despite the big sampling effort, mortality estimates could be underestimated since big percentages are butchered and discarded at sea. Still, numbers remains high with almost 1000 turtles in a five-year period and an illegal trade persists. Urgent measures are needed to recover this endangered species.

Keywords: *Chelonia mydas*, illegal captures, black market, Pisco, Perú.

The sea turtle consumption in the Pisco, San Andrés area (13°72'S, 76°22'W), ~130 km south of Lima has a long history. Records go back to the pre-hispanic era (4000-2000 AC), when around 30 turtles were reported from an archaeological site around Paracas (13°83'S, 76°25'W). During the Middle Ages, the Spanish Jesuit chronicler Bernabé Cobo reported the capture of ~90 turtles with a beach seine in Pisco (Frazier & Bonavia, 2000). Later, during the 1960s and 1970s, a report (Frazier, 1979) on the current state of the sea turtle fishery in Peru and the East Pacific, stated that the greatest numbers of turtles were captured in the port of Pisco, roughly 142 ton were landed, representing more than half the landings for the entire country from 1965-1976 period. But it is during late 1970s when a sophisticated turtle trafficking network thrived in the Pisco area. The fishery was operated by 7 to 10 boats exclusively dedicated to catch sea turtles, using nets specially designed to catch turtles, and with a well-established trafficking structure. The average catch was between 10 to 30 turtles per vessel per day and the captures were mainly composed by green turtles and, to a lesser extent, leatherbacks (Frazier, 1979; Hays-Brown & Brown, 1982). During the 1987 El Niño up to 110 boats were recorded landing turtles in Pisco (Zeballos, *pers. obs.*), standing as the largest sea turtle

harvest ever recorded in Peru. In just 10 months, over 20,000 turtles were landed (Aranda & Chandler, 1989). Sea turtle extraction was a legal activity until 1995 when a total ban on use of all the sea turtle species was established by Peruvian legislation (Morales & Vargas, 1996). However, the hunting still continues in Pisco by means of incidental and directed illegal captures as was preliminary evidenced (De Paz *et al.*, 2002, 2004, 2007; Quiñones *et al.*, 2010).

In order to collect evidence of sea turtle illegal captures in Pisco, the regional laboratory of the Peruvian Marine Research Institute (IMARPE) conducted an assessment. From November 2009 until March 2015, bi-weekly visits to local dumpsites and local beaches in the surrounding area of Pisco and San Andrés were conducted and all encountered turtle remains were registered and geo-referenced. The carcasses, carapaces and stranded individuals were identified to species level and the curved carapace length (CCL) was measured and registered. For black sea turtles (*Chelonia mydas agassizii*), life stages were characterized as follows: juvenile (<69 cm CCL), sub-adult (>69 cm and <85 cm CCL), and adult turtles (>85 cm CCL), according to the minimum breeding size (CCL = 69 cm) and mean breeding size (CCL = 85 cm) reported for 1037 nesting females in the Galapagos rookery during 2004-2007

(Zárate *et al.*, 2013). For olive ridleys (*Lepidochelys olivacea*), individuals >57 cm CCL were considered adults, using the minimum size reported for nesting females in Nancite, Costa Rica (Marquez, 1990). For leatherback sea turtles (*Dermochelys coriacea*), individuals >123 cm CCL were considered adults, using the minimum size reported for nesting females in Parque Nacional Marino las Baulas in Costa Rica (Reina *et al.*, 2002). In the case of hawksbill turtles, individuals >66 cm CCL were considered adults, based on the minimum size reported for nesting females at Estero Padre Ramos in Nicaragua (Altamirano *et al.*, 2014). All carapaces were painted with red spray to avoid recounting.

To identify the primary anthropogenic-derived threats for green turtles, the type of record obtained was used to determine the potential cause of death. If encountered at a dumpsite the turtle was considered to have been illegally captured, whereas if a turtle was found stranded at the beach the cause of death was classified either as boat collision, incidental entanglement in fisheries gear, or undetermined. Boat collisions were identified based on clear boat strike wounds or propeller injuries, whereas entanglements were considered when clear net derived injuries were present. We calculated annual mortality rates by dividing the number of new carcasses found at dumpsites (mean n° carcasses year⁻¹) and at the beach (mean n° carcasses km⁻¹ year⁻¹) by the time elapsed between surveys, using the methodology of Senko *et al.* (2014).

In addition, for each sea turtle reported, a decomposition stage was determined using a scale according to USGS standards (Work, 2000) which is: 1) Stranded dying animal; 2) fresh animal recently deceased; 3) animal moderately decomposed; 4) animal highly decomposed; and 5) skeletal or mummified animal. Necropsies were performed on fresh individuals in Category 2 and esophagus and stomachs were retrieved and immediately transported to the IMARPE regional laboratory in Pisco to analyze contents to determine diet composition. From stomach contents in black turtles, prey items were identified to the lowest possible taxon. Plant matter, mollusks, fish, crustaceans, jellyfish, and actinarian anemones were identified according to Dawson *et al.* (1964), Alamo & Valdivieso (1987), Chirichigno (1974), Retamal (1981), and Sanamyan *et al.* (2004). Quantitative assessment of diet was based on the relative wet weight (ww) in each sample, and was calculated as follows: %ww = (wet weight of a diet item/total weight of all items) × 100.

Finally, in order to determine the dimension of the black market and extent of illegal trade, we made several visits to the Pisco central market and conducted

70 structured qualitative interviews with local fishermen; main topics were by-catch and illegal trade. Between November 2009 and March 2015, five inland dumping places were reported, mainly in the central Pisco-San Andrés area. The majority of sea turtles (85%) were registered at the main dumping site (13°74'S, 76°20'W) and surrounding areas and the remainder (12.7%) of turtles were reported along the shorelines of Caucato, Pisco, San Andrés and Paracas (between 13°65'S and 13°86'S). A total of 953 sea turtles were registered: 92.2% (n = 898) were black turtles, 4.3% (n = 41) were olive ridleys, 1.4% (n = 13) were leatherbacks, and 0.1% (n = 1) were hawksbills. The mean CCLs were 59.2 ± 9.5 cm (range: 34.2-90.9 cm) for black turtles, 60.4 ± 6.2 cm CCL (range: 39.6-69.7 cm) for olive ridleys, 113 ± 18.2 cm CCL (range: 80-131.3 cm) for leatherbacks; the single juvenile hawksbill turtle had a CCL of 50.2 cm. Based on sizes observed, the life stage was determined: for black turtles, 83% were juveniles, 15.4% sub-adults and only 1.6% adults (Fig. 1), 38.5% of olive ridleys were juveniles, and 70% of leatherbacks were juveniles.

The main identified cause of death of black turtles was illegal capture (89%, n = 899). For this species, stranding events were represented by only 1.4% (n = 13), and within this 0.9% were not determined (n = 8), 0.3% were due to collisions (n = 3), and 0.2% due to entanglements (n = 2) (Fig. 1). For olive ridleys, 97.5% were illegal captures and only 2.5% were standings where the cause of death was not determined. For leatherbacks, 85% were illegal captures and 15% stranded. Of the stranded leatherbacks, half showed evidence of collision and the rest was not determined. Our mean black turtle mortality estimates were 8.1 carcasses km⁻¹ year⁻¹ at beaches and 160.2 carcasses year⁻¹ at dumpsites (Fig. 2).

According to the decomposition scale used, no dying animal was reported, 24.7% (n = 177) were classified as freshly deceased, 50.8% (n = 364) were moderately decomposed, 22.8% (n = 163) highly decomposed, and 1.7% (n = 12) skeletal or mummified. Five esophageal and stomach content analyses were performed during October 2014. The most consumed item, expressed in percentage of wet weight (% ww), was silverside fish (*Odontesthes regia regia*) eggs (47.9% ww), followed by the seaweed "yuyo" (*Chondracanthus chamissoi*, 31.4% ww) and the actinarian anemone *Paranthus* sp. (16.2% ww). Crustaceans, polychaetes, fish, and green seaweed remains were registered in lesser amounts (Table 1).

The existence of illegal trade was demonstrated. Recovered evidence at the Pisco Central fish-market observations during summer and autumn months indicates there were five ladies offering sea turtle meet




Figure 1. a) Size structure and causes of death of black turtles (*Chelonia mydas agassizii*), based in information from dumpsites and strandings in the Pisco area for the period 2009-2015, b) pie chart of main causes of death in black turtles, numbers represent the number of cases with their percentages.

Table 1. Prey wet weights (g) in both esophageal and stomach samples (n = 5) of black turtles (*Chelonia mydas*) and percentage of wet weight (%ww) in the Pisco area, during October 2014.

Prey item	1	2	3	4	5	Total	%ww
Animal matter							
Silverside eggs (fish)		181.3	4.5	580.9	55.0	821.7	47.9
<i>Paranthus</i> sp. (anemone)		70.3	88.2	37.6	82.1	278.2	16.2
Silverside (fish)			10.6			10.6	0.6
<i>Squilla stillirostris</i> (crustacean)			14.5		16.5	31.0	1.8
<i>Diopatra</i> (Polychaete)			1.2			1.2	0.1
Crustacean remains		0.5			3.7	4.2	0.2
Other Polychaetes					0.2	0.2	0.0
Vegetal matter							
<i>Chondracanthus chamissoi</i>	12.3	95.9	210.0	67.6	153.3	539.0	31.4
<i>Enteromorpha oficinalis</i>	16.4					16.4	1.0
<i>Ulva papenfusi</i>	6.5					6.5	0.4
<i>Caulerpa filiformis</i>	2.8		3.6			6.4	0.4
Mineral matter							
Mud	0.8					0.8	0.05

for sale. Each one had approximately two turtles for sale and the price per kg varied from US\$ 7 to 10. Based on our observations, we infer that between 10 to 20 turtles, mainly black turtles, are sold per week during summer and autumn months, whereas during winter and early spring season the amounts of turtle for sale were less and some weeks were totally absent from the market. As a result of the interviews, the illegal turtle trade was divided into three main activities: i) Local consumption, subdivided in family consumption by fishermen and subsistence consumption on board fishing vessels by fishermen; ii) market and other types of commercialization, subdivided in turtle meat and fins sold in the Pisco market as well as door to door sales of turtle meat in the fishing town of San Andrés;

and iii) demand from other towns, subdivided in demands from Lima, the capital city, and demands from Pisco nearby towns like Chinchá, Cañete, and Ica. In addition, as a result of combining answers from interviews and night observations, two fishermen, pretending to catch silverside fish at Paracas, were observed landing turtles hidden inside large plastic bags. These fishermen had artisanal gillnets with a mesh size of 60 cm (the traditional measure for turtle nets). The majority of the prey items consumed by the turtles captured were animal matter, mainly represented by silverside fish eggs and anemones (*Paranthus* sp.) during spring, whereas during summer months the diet was mainly composed of jellyfish. There was a clear seasonality in prey consumption, with jellyfish *Chrysaora*



Figure 2. Pictures of illegal captures of sea turtles in the Pisco area. Clockwise from the first picture top left: leatherback captured for human consumption at the beach; and the three remaining ones were black turtles in the main dumpsite of Pisco.

plocamia consumption occurring mainly during the summer-autumn seasons and sea anemone *Paranthus* sp. consumption during the winter-spring seasons (Paredes, 2015).

The human activity that has the largest impact on sea turtles is fisheries bycatch (Lewinson *et al.*, 2004, 2013; Wallace *et al.*, 2011), even worse, direct harvest remains a major threat to sea turtles population worldwide (Campbell, 2003; Lewison *et al.*, 2013), as was previously stated a legal sea turtle fishery existed in Pisco until mid-90s, after that illegal poaching continues in the country, however sea turtles poaching does not occurred exclusively in Peru, in several

countries in the Americas this illegal activity still occurs nowadays, like in the Mexican Pacific (Mancini & Koch, 2009; Mancini *et al.*, 2011; Senko *et al.*, 2014), the wide Caribbean area (Bell *et al.*, 2006; Richardson *et al.*, 2009; Lagueux *et al.*, 2014) and Brazil (Geubert *et al.*, 2013), in addition, this illegal activity occurs elsewhere worldwide, like in Africa (Riskas & Tiwari, 2013); Southeast Asia (Joseph *et al.*, 2014); Oceania (Maison *et al.*, 2010) among other places.

Direct extraction, together with by-catch interaction with small scale gillnets fisheries (SSGF), is a strong factor of mortality in Peru mostly due to economic

needs, lack of law enforcement and strong traditional consumption, as was reported in BC, Mexico (Mancini *et al.*, 2011). The SSGF usually operates in nations where there are few protective measures and limited enforcement capabilities (Chuenpagdee *et al.*, 2006; Dutton & Squires, 2008). Peru is a typical example of this situation, where an estimated of ~6000 turtles are extracted annually by SSGF (Alfaro-Shigueto *et al.*, 2011), particularly in the Pisco area, where still nowadays fishermen are manufacturing “redes tortugeras” a traditional gill net for direct turtle harvest. In addition, retaining bycatch is a common practice in the area. Some artisanal fishermen go at sea with a wide array of legal nets; however, “redes tortugeras” are also taken and placed in hidden places on board, in order to use it “just in case” the target capture is not enough.

Despite the high sampling effort, the mortality estimates presented here are underestimated for at least two reasons. First, poachers are killing turtles at sea and throwing the carcasses into the water where they go unnoticed, similar to reports from BC, Mexico (Mancini & Koch, 2009). Second, the remains could be burned between the sampling days, thereby not allowing us to find the carcasses. Yet despite this, almost 1000 carapaces were encountered in the Pisco area. The high mortality rates of 160 turtles year⁻¹ in dumpsites and 8.1 km⁻¹ year⁻¹ at beaches are the consequence of a strong sampling effort (two times per week), a good spatial coverage (almost 100% of dumpsites in Pisco), and sea turtle aggregations in Pisco restricted to a small geographic area of less than 2.500 m² and ~200 km of coastline (Velez-Zuazo *et al.*, 2014; Quiñones *et al.*, 2017). It is suggested that mortality rates in Pisco are likely underestimated considering that this area overlaps and interacts with a strong SSGF. There are an average of 200 SSGF boats operating in San Andrés and 100 boats in Tambo de Mora.

The majority of the turtles illegally taken in Peru were small individuals, recruited from oceanic epipelagic areas to neritic areas (Luschi *et al.*, 2003) like in the shallow waters of Paracas Bay, indeed, our black turtle size structure reflected a predominance of juveniles (over 80%), however if we compared it with information from 1987, a size decrease of ~9 cm of CCL was noticed, in marine resources fisheries population assessments revealed that age and sexual maturation (ASM) could decrease as a consequence of fishing pressure (Olsen *et al.*, 2004; Swain *et al.*, 2007), in the case of sea turtles, a synergy of different issues like: fishing pressure, physical and biological characteristics of the environment and density dependent factors could influence the ASM and the growth rates of sea turtles (Avens & Snover, 2013).

Regarding the black market trade, the situation seems to be declining since the total ban in 1996 (Morales & Vargas, 1996). Nevertheless, we found that there are still local/regional markets that supply sea turtle meat and derived products. Though a quite strong regional trade persists in the area and this activity represents an extra income to fishermen in Pisco, where a turtle of average weight (30 kg) can be sold up to US\$60 demands of some turtle derived products persist, for instance, oil and turtle blood were thought to be effective remedies against flu and other diseases in the past, even nowadays the turtle oil is still demanded by local people, each bottle of 500 mL is sold at ~US\$10, instead turtle blood ingesting is no longer practiced. The trafficking structure fluctuates seasonally with a well-structured network during summer and autumn seasons and a weak and unstructured network during winter and spring seasons, however the current demand is much lower than in the former years.

All these illegal activities are prone to continue unless localized law enforcement and political measures are strongly implemented in the area. Even if our data is only a fraction of the whole illegal captures and trade in the country, it provides a valuable understanding of the current situation in the Pisco area. The presented results could be useful for further management and recommendation actions.

REFERENCES

- Alamo, V. & V. Valdivieso. 1987. Lista sistemática de moluscos marinos del Perú. Boletín Instituto del Mar del Perú & Deutsche Gesellschaft für Technische Zusammenarbeit, Volumen Extraordinario, Callao, Perú: 1-205.
- Alfaro-Shigueto, J., J.C. Mangel, F. Bernedo, P.H. Dutton, J.A. Seminoff & B.J. Godley. 2011. Small-scale fisheries of Peru: a major sink for marine turtles in the Pacific. *J. Appl. Ecol.*, 48: 1432-1440.
- Altamirano, E., V. Gadea, A. Gaos & M. Liles. 2014. Protocolo de monitoreo, protección de nidos y manejo de vivero de tortuga Carey (*Eretmochelys imbricata*) en la RN Estero Padre Ramos y Aserradores, Temporada 2014. Fauna y Flora Internacional e ICAPO, 34 pp.
- Aranda, C.A. & M.W. Chandler. 1989. Las tortugas marinas del Perú y su situación actual. *Bol. Lima*, 11: 77-86.
- Avens, L. & M.L. Snover. 2013. Age and age estimation in sea turtles. In: J. Wyneken, J. Lohmann & J.A. Musick (eds.). *The biology of sea turtles*. CRC Press, Boca Raton, pp. 3-97.

- Bell, C.D., J.M. Blumenthal, T.J. Austin, J.L. Solomon, G. Ebanks-Petrie, A. Broderick & B.J. Godley. 2006. Traditional Caymanian fishery may impede local marine turtle population recovery. *Endanger. Species Res.* 2: 63-69.
- Campbell, L.M. 2003. Contemporary culture, use, and conservation of sea turtles. In: P.L. Lutz, J.A. Musick & J. Wyneken (eds.). *Biology of sea turtles*. CRC, Boca Raton, pp. 307-338.
- Chirichigno, N. 1974. Clave para identificar los peces marinos del Perú. *Informe Instituto del Mar del Perú, Callao*, 44: 1-388.
- Chuenpagdee, R., L. Liguori, M.L.D. Palomares & D. Pauly. 2006. Bottom up, global estimates of small-scale fisheries catches. *Fisheries Centre Research Report, British Columbia University*, 14: 105 pp.
- Dawson, E.Y., C. Acleto & N. Foldvik. 1964. The seaweeds of Perú. *Nova Hedwigia*, 13: 1-111.
- De Paz, N., J.C. Reyes & M. Echegaray. 2002. Datos sobre captura, comercio y biología de tortugas marinas en el área de Pisco-Paracas. *Memorias I Jornada Científica Reserva Nacional de Paracas, Universidad Nacional Agraria La Molina, Lima*, pp. 125-129.
- De Paz, N., J.C. Reyes & M. Echegaray. 2004. Capture and trade of marine turtles at San Andrés, Southern Peru. In M.S. Coyne & R.D. Clark (eds.). *Proceedings of the Twenty-First Annual Symposium on Sea Turtle Biology and Conservation*. NOAA Technical Memorandum NMFS-SEFSC, 528: 52-54.
- De Paz, N., J.C. Reyes, M. Echegaray, M. Ormeño & H. Anchante. 2007. Identificación y manejo de hábitats críticos de tortugas marinas en Perú: Paracas, estudio de caso. In: C. Guerra-Correa, A. Fallabrino, P. Bolados-Díaz & C. Turner (eds.). *Estado actual y perspectivas de la investigación y conservación de las tortugas marinas en las costas del Pacífico Sur-Oriental*. Antofagasta, 28 pp.
- Dutton, P.H. & D. Squires. 2008. Reconciling biodiversity with fishing: a holistic strategy for Pacific sea turtle recovery. *Ocean Develop. Int. Law*, 39: 200-222.
- Frazier, J. 1979. Marine turtles in Peru and the east Pacific. Office of Zoological Research, National Zoological Park, Smithsonian Institution (unpublished manuscript).
- Frazier, J. & D. Bonavia. 2000. Prehispanic marine turtles in Peru: where were they? In: F.A. Abreu-Grobois, R. Briceño-Dueñas, R. Márquez & L. Sarti (eds.). *Proceedings of the Eighth International Sea Turtle Symposium*. U.S. Department Commerce, NOAA Tech. Memo NMFS-SEFSC-436: 243-245.
- Geubert, F.M., M. Barletta & M.F. da Costa. 2013. Threats to sea turtle populations in the Western Atlantic: poaching and mortality in small-scale fishery gears. *J. Coast. Res.*, 65: 42-47.
- Hays-Brown, C. & W.M. Brown. 1982. Status of sea turtles in the southeastern Pacific: emphasis on Peru. In: K.A. Bjorndal (ed.). *Biology and conservation of sea turtles*. Smithsonian Institution Press, Washington DC, pp. 235-240.
- Joseph, J., C.Y. Kuen, P.M. Palaniappan & L.H. Chark. 2014. Genetic investigation of green turtles (*Chelonia mydas*) harvested from a foraging ground at Mantanani, Sabah, Malaysia. *Herpetol. Conserv. Biol.*, 3: 516-523.
- Lagueux, C.J., C.L. Campbell & S. Strindberg. 2014. Artisanal green turtle, *Chelonia mydas*, fishery of Caribbean Nicaragua: I. Catch rates and trends, 1991-2011. *PLoS ONE*, 9: e94667.
- Lewison, R.L., L.B. Crowder, A.J. Read & S.A. Freeman. 2004. Understanding impacts of fisheries bycatch on marine megafauna. *Trends Ecol. Evol.*, 19: 598-604.
- Lewison, R.L., B. Wallace, J. Alfaro-Shigueto, J.C. Mangel, S.M. Maxwell & E. Hazen. 2013. Fisheries bycatch of marine turtles, lessons learned from decades of research and conservation. In: J. Wyneken, K.J. Lohmann & J.A. Musick (eds.). *The biology of sea turtles*. Vol. 3. CRC, Boca Raton, pp. 329-351.
- Luschi, P., G.C. Hays & F. Papi. 2003. A review of long-distance movements by marine turtles, and the possible role of ocean currents. *Oikos*, 103: 293-302.
- Maison, K.A., I.K. Kelly & K.P. Frutchey. 2010. Green turtle nesting sites and sea turtle legislation throughout Oceania. US Department of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service. NOAA Technical Memorandum NMFS-F/SPO-110.
- Mancini, A. & V. Koch. 2009. Sea turtle consumption and black market trade in Baja California Sur, Mexico. *Endanger. Species Res.*, 7: 1-10.
- Mancini, A., J. Senko, R. Borquez-Reyes, J.G. Póo, J.A. Seminoff & V. Koch. 2011. To poach or not to poach an endangered species: elucidating the economic and social drivers behind illegal sea turtle hunting in Baja California Sur, Mexico. *Hum. Ecol.*, 39: 743-756.
- Márquez, R. 1990. Sea turtles of the world. An annotated and illustrated catalogue of sea turtle species known to date. *FAO Fish. Synop. No. 125, Vol. 11, Rome*, 81 pp.
- Morales, V.R. & P. Vargas. 1996. Legislation protecting marine turtles in Peru. *Mar. Turtle Newslett.*, 75: 22-23.
- Olsen, E.M., H. Knutsen, J. Gjøsaeter, P.E. Jorde, J.A. Knutsen & N. Stenseth. 2004. Life-history variation among local populations of Atlantic cod from the Norwegian Skagerrak coast. *J. Fish. Biol.*, 64: 1725-1730.

- Paredes, E. 2015. Hábitos alimentarios de la tortuga verde del Pacífico este *Chelonia mydas agassizii* (Boucort, 1868) en la bahía de Paracas, Ica, Perú, durante el año 2010. Tesis Universidad Nacional Mayor de San Marcos, San Marcos, 61 pp.
- Quiñones, J., S. Quispe, E. Paredes. 2017. Black turtle population dynamics in Paracas, Peru during a six year in water monitoring. Proceedings of the Thirty Six Annual Symposium on Sea Turtle Biology and Conservation. Lima, Perú, March 2016. (in press).
- Quiñones, J., V.G. Carman, J. Zeballos, S. Purca & H. Mianzan. 2010. Effects of El Niño-driven environmental variability on black turtle migration to Peruvian foraging grounds. *Hydrobiologia*, 645: 69-79.
- Reina, R.D., P.A. Mayor, J.R. Spotila, R. Piedra & F.V. Paladino. 2002. Nesting ecology of the leatherback turtle, *Dermochelys coriacea*, at Parque Nacional Marino Las Baulas, Costa Rica: 1988-1989 to 1999-2000. *Copeia*, 2002: 653-664.
- Retamal, M.A. 1981. Catálogo ilustrado de los crustáceos decápodos de Chile. Universidad de Concepción, Gayana, 44: 1-110.
- Richardson, P.B., M.W. Bruford, M.C. Calosso, L.M. Campbell, W. Clerveaux, A. Formia, B.J. Godley, A.C. Henderson, K. McClellan, S. Newman, K. Parsons, M. Pepper, S. Ranger, J. Silver, L. Slade & A. Broderick. 2009. Marine turtles in the Turks and Caicos Islands: remnant rookeries, regionally significant foraging stocks, and a major turtle fishery. *Chelonian Conserv. Biol.*, 8: 192-207.
- Riskas, K.A. & M. Tiwari. 2013. An overview of fisheries and sea turtle bycatch along the Atlantic coast of Africa. *Munibe*, 1: 1-82.
- Sanamyan, N., K. Sanamyan & D. Schories. 2004. Actiniaria.com. Available in [<http://actiniaria.com>]. Reviewed: 18 January 2017.
- Senko, J., A. Mancini, J.A. Seminoff & V. Koch. 2014. By-catch and directed harvest drive high green turtle mortality at Baja California Sur, Mexico. *Biol. Conserv.*, 169: 24-30.
- Swain, D.P., S.F. Sinclair & J.M. Hanson. 2007. Evolutionary response to size-selective mortality in an exploited fish population. *Proc. R. Soc. Lond. B*, 274: 1015-1022.
- Velez-Zuazo, X., J. Quiñones, A.S. Pacheco, L. Klinge, E. Paredes, S. Quispe & S. Kelez. 2014. Fast growing, healthy and resident green turtles (*Chelonia mydas*) at two neritic sites in the central and northern coast of Peru: implications for conservation. *PLoS ONE*, 9: e113068.
- Wallace, B.P., A.D. DiMatteo, A.B. Bolten, M.Y. Chaloupka, B.J. Hutchinson, F.A. Abreu-Grobois, J.A. Mortimer *et al.* 2011. Global conservation priorities for marine turtles. *PLoS ONE*, 6:e24510.
- Work, T.M. 2000. Sea turtle necropsy manual for biologists in remote refuges. National Wildlife Health Center, Hawaii Field Station, 25 pp.
- Zárate, P., K.A. Bjorndal, M. Parra, P.H. Dutton, J.A. Seminoff & A.B. Bolten. 2013. Hatching and emergence success in green turtle *Chelonia mydas* nests in the Galápagos Islands. *Aquat. Biol.*, 19: 217-229.

Received: 30 August 2016; Accepted: 13 January 2017