

Short Communication

New record and morphometry of the bluntnose sixgill shark *Hexanchus griseus* (Chondrichthyes: Hexanchidae) in Baja California Sur, Mexico

**Edgar Eduardo Becerril-García^{1,3}, Carlos Augusto Aguilar-Cruz²
Alexis Alejandro Jiménez-Pérez² & Felipe Galván-Magaña¹**

¹Instituto Politécnico Nacional, Centro Interdisciplinario de Ciencias Marinas, La Paz, BCS, México

²Universidad Autónoma de Baja California Sur, La Paz, BCS, México

³Pelágicos Kakunjá A.C., La Paz, BCS, México

Corresponding author: Felipe Galván-Magaña (galvan.felipe@gmail.com)

ABSTRACT. We report the first record and morphometry of a complete juvenile *Hexanchus griseus* female shark, caught in Punta Lobos, Baja California Sur, Mexico. The oceanographic conditions of the area, where the specimen was caught, coincide with the habitat previously described for this shark species, with low temperature, stable salinity and high hydrostatic pressure. There is an overall lack of information about *H. griseus* in Mexico, with only eight records of its presence in the Mexican Pacific. Moreover, none of these previous records provided detailed information on the morphometry of a whole bluntnose sixgill shark. Here, we present the morphometric measures of a complete individual caught in shallow waters. We expect this information to be useful for future comparisons of specimens caught in other coastal regions.

Keywords: *Hexanchus griseus*, elasmobranchs, distribution, bathyal zone, Tropical Eastern Pacific.

The bluntnose sixgill shark, *Hexanchus griseus*, is the only species of the genus *Hexanchus* recorded in the Mexican Pacific (Compagno, 1984; Fischer *et al.*, 1995). It is one of the largest carnivorous sharks, occupying the highest trophic level, reaching 4.82 m in total length (Ebert, 2015). The species possesses six pairs of gill slits, a subterminal mouth and a short, broadly rounded blunt snout. It is grey with or without irregular brown spots. Also, the distance from the dorsal fin to the caudal fin is equal to the length of the dorsal fin base (Compagno, 1984).

Records of *H. griseus* in Mexico have come from catches and fish lists drawn up in Sonora (Santana-Morales, 2005; Robertson & Allen, 2015), Baja California (Miller & Lea, 1972; Castro-Aguirre & García-Domínguez, 1988; Goodson, 1988; Ruiz-Campos *et al.*, 2010) and Baja California Sur (Castro-Aguirre *et al.*, 2003; Robertson & Allen, 2015). Of the studies reporting the occurrence of this species, that of Castro-Aguirre *et al.* (2003) is the only one, however, that includes morphometric measurements, those of the head of an unsexed specimen recorded in Bahía Magdalena, Baja California Sur, Mexico. Here, we present the first record and morphometry of a complete

specimen of 107 cm in total length, caught 180 km south of Bahía Magdalena, off the fishing camp of Punta Lobos, Baja California Sur, Mexico.

A juvenile female of *H. griseus* was caught incidentally on 18 December 2013 by coastal fishermen using longlines. Punta Lobos is located at 23°25'N, 110°14'W, on the west coast of the Baja California Peninsula (Fig. 1).

In this area, oceanographic conditions at a depth of 200 to 2500 m are characterized by a temperature range of 4 to 12°C, a stable salinity of 34 to 35, and dissolved oxygen values of 2.5 to 4 mL L⁻¹ (Locarnini *et al.*, 2013; Zweng *et al.*, 2013; García *et al.*, 2014). Such conditions coincide with the known habitat of *H. griseus* in its area of distribution, *i.e.*, hypoxic environments with low temperature and high hydrostatic pressure (Compagno, 1984; Barnett *et al.*, 2012). Although this species mainly inhabits the continental slope, juveniles have been reported to make incursions into neritic environments (over the continental shelf) during the night, when they undergo vertical migration possibly related to feeding (Compagno, 1984; Andrews *et al.*, 2007, 2009, 2010; Comfort, 2012).




Figure 1. Previous records of *Hexanchus griseus* and new location of Punta Lobos, BCS, Mexico.

Table 1. Morphometry of the bluntnose sixgill shark female captured off Punta Lobos, BCS, Mexico.

Measure	Length (cm)	Proportion (%)
TL (total length)	107.2	100.00
FL (fork length)	84.6	78.92
PCL (pre-caudal-fin length)	74.9	69.87
HDL (head length)	22.1	20.62
POB (preorbital length)	5.2	4.85
PP1 (prepectoral-fin length)	22.1	20.62
PP2 (prepelvic-fin length)	27.8	25.93
PAL (preanal-fin length)	63.4	59.14
EYL (eye length)	1.6	1.49
ING (intergill length)	1.4	1.31
P1A (pectoral-fin anterior margin)	5.4	5.04
P1B (pectoral-fin base)	15.3	14.27
CDM (dorsal caudal-fin margin)	10.3	9.61
CPV (oreventral caudal-fin margin)	30.1	28.08
CFW (caudal-fin fork width)	3.4	3.17
CFL (caudal-fin fork length)	5.4	5.04
P2L (pelvic-fin length)	2.9	2.71
P2B (pelvic-fin base)	8.7	8.12
ANL (anal-fin length)	1.7	1.59
ANA (anal-fin anterior margin)	8.2	7.65
ANB (anal-fin base)	5.4	5.04

ry of Fish Ecology of the Interdisciplinary Centre of Marine Sciences (CICIMAR-IPN, by its Spanish acronym) in La Paz, BCS, Mexico.

Despite the fact that *H. griseus* is a top predator, knowledge of this shark species is scarce and insufficient (Ebert, 2002, 2003), in both coastal and



Figure 2. Juvenile female of *Hexanchus griseus* caught off the fishing camp of Punta Lobos, BCS, Mexico.

The caught specimen was dark grey without spots on the back and white on the belly, with no clear boundary between the two colours (Fig. 2). Its flabby body, typical of organisms inhabiting the mesopelagic or bathypelagic zone (Randall & Farrell, 1997), had an epidermis entirely covered with mucus, with no ectoparasites detected.

Taxonomic identification was performed using Fischer *et al.* (1995) dichotomous key as well as morphometry of the whole organism (Table 1). The shark specimen was transported on ice to the Laborato-

deep-water ecosystems (Barnett *et al.*, 2012). This study provides the first record of *H. griseus* that includes detailed morphometric data of a complete individual. This individual was caught in shallow waters off the west coast of Baja California Sur, providing additional information on the species' distribution

in Mexico. As for its conservation status, the bluntnose sixgill shark is categorised as “near threatened” by the IUCN (Cook & Compagno, 2005).

In Mexico, information relative to *H. griseus* is insufficient for realize a proper assessment of the status of its population; furthermore, the limited information on its reproductive biology suggests that this taxon may be vulnerable to fishing (Barnett *et al.*, 2012). The punctual catches recording of *H. griseus* taking advantage of them to produce scientific knowledge about basic aspects of its biology, will allow better understanding of the ecological niche occupied by this species in the national territory.

ACKNOWLEDGEMENTS

We thank the fishermen of Punta Lobos, and Nefertiti Taydé Roldán-Wong, and Janna Domínguez-Boza. Nicole Reguera-Rouzaud for her help in drawing the figures. FGM thanks the Instituto Politecnico Nacional for funding from the Commission of Operation and Promotion of Academic Activities (COFAA) and Performance Incentive for Researchers (EDI) grants.

REFERENCES

- Andrews, K.S., G.D. Williams & P.S. Levin. 2010. Seasonal and ontogenetic changes in movement patterns of sixgill sharks. *PLoS ONE*, 5: e12549.
- Andrews, K.S., P.S. Levin, S.L. Katz, D. Farrer, V.F. Galluci & G. Bargmann. 2007. Acoustic monitoring of sixgill shark movements in Puget Sound: evidence for localized movement. *Can. J. Zool.*, 85: 1136-1142.
- Andrews, K.S., G.D. Williams, D. Farrer, N. Tolimieri, C.J. Harvey, G. Bargmann & P.S. Levin. 2009. Diel activity patterns of sixgill sharks, *Hexanchus griseus*: the ups and downs of an apex predator. *Anim. Behav.*, 78: 525-536.
- Barnett, A., J.M. Braccini, C.A. Awruch & D.A. Ebert. 2012. An overview on the role of Hexanchiformes in marine ecosystems: biology, ecology and conservation status of a primitive order of modern sharks. *J. Fish Biol.*, 80: 966-990.
- Castro-Aguirre, J.L. & F. García-Domínguez. 1988. Sobre la presencia de *Isistius brasiliensis* (Quoy y Gaimard) (Squaliformes: Squalidae: Dalatiinae) en el Golfo de México, con un elenco sistemático de las especies mexicanas pertenecientes al Superorden Squalomorphii. *An. Esc. Nac. Cienc. Biol.*, 32: 91-108.
- Castro-Aguirre, J.L., A. Antuna-Mendiola, A.F. González-Acosta & J. De La Cruz-Agüero. 2003. Primer registro del tiburón cañabota *Hexanchus griseus* (Bonnaterre, 1788) (Chondrichthyes: Selachii-morpha) en la costa occidental de Baja California Sur, México. *CICIMAR Oceánides*, 18(1): 39-41.
- Compagno, L.J.V. 1984. Sharks of the world. An annotated and illustrated catalogue of shark species known to date. Part 1. Hexanchiformes to Lamniformes. FAO species catalogue, Rome, 125(4): 1-249.
- Comfort, C. 2012. Spatial and trophic ecology of the bluntnose sixgill shark: environmental drivers of behavior and comparative trophic position in two distinct habitats. Master’s Thesis, University of Hawaii Manoa, Honolulu, 114 pp.
- Cook, S.F. & L.J.V. Compagno. 2005. *Hexanchus griseus*. The IUCN Red List of Threatened Species 2005. [<http://dx.doi.org/10.2305/IUCN.UK.2005.RLTS.T10030A3155348.en>]. Reviewed: 17 August 2016.
- Ebert, D.A. 2002. Some observations on the reproductive biology of the sixgill shark *Hexanchus griseus* (Bonnaterre, 1788) from southern African waters. *S. Afr. J. Mar. Sci.*, 24: 359-363.
- Ebert, D.A. 2003. Sharks, rays and chimaeras of California. University of California Press, Berkeley, 284 pp.
- Ebert, D.A. 2015. Deep-sea cartilaginous fishes of the southeastern Atlantic Ocean. FAO, Rome, 251 pp.
- Fischer, W., F. Krupp, W. Schneider, C. Sommer, K.E. Carpenter & V.H. Niem. 1995. Guía FAO para la identificación de especies para los fines de la pesca. Pacífico centro-oriental. Volumen II. Vertebrados-Parte 1, FAO, Rome, 564 pp.
- García, H.E., R.A. Locarnini, T.P. Boyer, J.I. Antonov, O.K. Baranova, M.M. Zweng, J.R. Reagan & D.R. Johnson. 2014. World Ocean Atlas 2013. In: S. Levitus & A. Mishonov (eds.). Dissolved oxygen, apparent oxygen utilization, and oxygen saturation. NOAA Atlas NESDIS 75, NOAA, 27 pp.
- Goodson, G. 1988. Fishes of the Pacific coast: Alaska to Peru, including the Gulf of California and the Galapagos Islands. Stanford University Press, Stanford, 267 pp.
- Locarnini, R.A., A.V. Mishonov, J.I. Antonov, T.P. Boyer, H.E. Garcia, O.K. Baranova, M.M. Zweng, C.R. Paver, J.R. Reagan, D.R. Johnson, M. Hamilton & D. Seidov. 2013. World Ocean Atlas 2013, Temperature. In: S. Levitus & A. Mishonov (eds.). NOAA Atlas NESDIS 73, NOAA, 40 pp.
- Miller, D.J. & R.N. Lea. 1972. Guide to the coastal marine fishes of California. Fish. B-NOAA, 157: 1-249.
- Randall, D.J. & A.P. Farrell. 1997. Deep-sea fishes. Academic Press, San Diego, 388 pp.
- Robertson, D.R. & G.R. Allen. 2015. Peces costeros del Pacífico Oriental Tropical: sistema de información en

- Línea. [<http://biogeodb.stri.si.edu/sftep/es/pages>]. Reviewed: 30 August 2016.
- Ruiz-Campos G., J.L. Castro-Aguirre, E.F. Balart, L. Campos-Dávila & R. Vélez-Marín. 2010. New specimens and records of chondrichthyan fishes (Vertebrata: Chondrichthyes) off the Mexican Pacific coast. *Rev. Mex. Biodivers.*, 81: 363-371.
- Santana-Morales, O. 2005. Catálogo de tiburones, rayas y quimeras (Pisces Chondrichthyes) que habitan en las aguas del norte del Golfo de California. Tesis de Licenciatura. Universidad de Guadalajara, Guadalajara, 108 pp.
- Zweng, M.M, J.R. Reagan, J.I. Antonov, R.A. Locarnini, A.V. Mishonov, T.P. Boyer, H.E. Garcia, O.K. Baranova, D.R. Johnson, D. Seidov & M.M. Biddle. 2013. World Ocean Atlas 2013, Salinity. In: S. Levitus & A. Mishonov (eds.). Technical NOAA Atlas NESDIS, 74: 39 pp.

Received: 20 September 2016; Accepted: 24 January 2017