

Research Article

Spatial variation in biometric parameters and condition index of *Modiolus capax* in contrasting coastal environments in the Gulf of California

Fernando A. Aguilar-Santana¹ , Alma R. Rivera-Camacho¹ 

Ilie S. Racotta²  & Marcial Arellano-Martínez¹ 

¹Instituto Politécnico Nacional, Centro Interdisciplinario de Ciencias Marinas (CICIMAR)
La Paz, Baja California Sur, Mexico

²Centro de Investigaciones Biológicas del Noroeste, S.C. (CIBNOR), La Paz
Baja California Sur, Mexico

Corresponding author: Fernando A. Aguilar-Santana (fagsa.03@gmail.com)
Marcial Arellano-Martínez (marellam@ipn.mx)

ABSTRACT. The biometric parameters and organismal condition of marine bivalves are strongly influenced by environmental conditions such as temperature, food availability, and anthropogenic disturbance. Using biometric parameters, the condition index (CI), and weight-length relationships (WLR), this study evaluated spatial variation in the organismal condition of *Modiolus capax* at coastal sites with contrasting environmental conditions in the Gulf of California, Mexico. Mussels from the Santa Rosalía (SR) mining port, a site historically impacted by mining activities and characterized by lower temperatures and higher food availability (chlorophyll-*a*), exhibited larger shell sizes and greater visceral mass, along with negative allometric growth patterns associated with shell elongation. In contrast, mussels from the La Paz Bay sites with higher temperatures and lower food availability, and with different types and intensities of anthropogenic influence (thermoelectric plant and phosphorite mine), were consistently smaller and lighter. Biometric parameters, CI, and WLR revealed clear spatial variation, indicating marked differences in morphological responses and somatic growth patterns among sites. However, at SR, shell elongation altered growth allometry and skewed condition-index interpretations based on shell length, artificially lowering CI values despite high visceral mass. These differences indicate that local environmental conditions strongly modulate biometric parameters across sites and suggest that morphological responses may mask organismal condition when conventional indices are applied without accounting for shell shape variation associated with potentially stressful conditions. Overall, this study highlights the value and limitations of biometric and condition indices as integrative indicators of population dynamics, providing a baseline to integrate future biometric assessments with physiological and biochemical biomarkers in historically impacted coastal systems.

Keywords: *Modiolus capax*; environmental heterogeneity; anthropogenic impacts; morphological plasticity; allometric growth, Mexico

INTRODUCTION

In marine ecosystems, environmental variation influences the dynamics that govern ecosystem structure and functioning (Doney et al. 2012, Assan et al. 2020).

These variations create favorable conditions for some species or populations, such as optimal temperatures and greater food or nutrient availability, which optimize physiological regulation and reduce competition and predation (Johan et al. 2015, Canosa & Bertucci

2023). However, when environmental conditions fluctuate beyond the normal range of variability, the resulting stress causes suboptimal physiological performance, leading to adverse consequences, such as decreased growth, smaller body size, lower reproductive capacity, and, in extreme cases, increased mortality (Doney et al. 2012, Johan et al. 2015). Therefore, anthropogenic impacts substantially alter organisms' natural habitats, exacerbating environmental stress (Brander et al. 2010, Moullec et al. 2021). These impacts, combined with low food availability and abrupt temperature changes outside the optimal range, significantly deteriorate marine populations (Assan et al. 2020, Horwitz et al. 2020). However, some species have developed adaptation mechanisms that allow them to tolerate extreme environmental conditions (Przeslawski et al. 2005), as do intertidal bivalves, which inhabit the intertidal zone, one of the most stressful habitats (Gosselin & Chia 1995). For this reason, bivalves are considered resilient to fluctuating environmental conditions (Chelyadina et al. 2022, Sampaio et al. 2022, Sepúlveda et al. 2024), so studying them allows detection of adverse impacts before they become irreversible (Helmholz et al. 2016).

Somatic indices, biometric parameters, and weight-length relationships (WLR) are key diagnostic tools for assessing organisms' physiological health in response to environmental changes. Their sensitivity to habitat conditions and their capacity to reflect essential physiological processes (growth, reproduction, and others) in organisms (Mercado-Silva 2005, Ramesha & Sophia 2015) make them essential methodological approaches in ecotoxicology, conservation, and management of aquatic resources since they provide information on stressful conditions caused by environmental factors and the effects of pollution on biota (Lucas & Beninger 1985, Filgueira et al. 2013).

Mussels, particularly species belonging to the genera *Mytilus* and *Modiolus*, are known for their high tolerance to stress conditions, such as low food availability, temperature fluctuations, and pollution (Anestis et al. 2008, Muñoz-Barbosa & Huerta-Díaz 2013, Clark et al. 2021). These bivalves are widely distributed in the Gulf of California (GC), a region of high biological, oceanographic, and geochemical relevance, recognized for its elevated primary productivity and diversity of marine habitats (Álvarez-Borrego 2010, Páez-Osuna et al. 2017). The GC has remarkable environmental heterogeneity driven by spatial and temporal variability in temperature, oxygen concentration, salinity, and nutrient availability, which promotes habitat and microhabitat formation and

shapes the structure and species composition of biological communities (Brusca 2010, Brusca & Hendrickx 2010). Additionally, several areas of the GC have been impacted by anthropogenic activities, including mining and metal ore processing, industrial discharges, municipal effluents, and thermoelectrical plants (Páez-Osuna et al. 2017). Among the benthic communities inhabiting this region is the mussel *Modiolus capax*, a species of local commercial and subsistence importance (local fishermen, *pers. comm.*). The biology of *M. capax* has been extensively studied in the GC, with research addressing its reproductive biology, recruitment patterns, capacity to bioaccumulate pollutants, and environmentally driven morphological variability (Gutiérrez-Galindo et al. 1999, Muñoz-Barbosa & Huerta-Díaz 2013, López-Carvalho et al. 2017, García-Corona et al. 2018, García-Domínguez et al. 2018, Roldán-Wong et al. 2020, Góngora-Gómez et al. 2022). However, studies evaluating the combined effects of environmental variability and pollutant exposure on *M. capax* biological responses along the western coast of the GC remain limited.

By assessing biometric parameters, condition index (CI), and WLR, this study evaluated spatial variation in the physiological condition of *M. capax* across contrasting coastal environments driven primarily by differences in temperature, food availability, and pollutant exposure. The research was conducted at four sites along the western coast of the GC: the mining port of Santa Rosalía (SR), characterized by elevated metal concentrations in sediments, lower water temperatures, and higher food availability, and three sites within La Paz Bay (LPB), characterized by higher temperatures, lower food availability, and different types and intensities of anthropogenic influence.

MATERIAL AND METHODS

Study area

The study was conducted at four sites within the GC (Fig. 1) between 2017 and 2020. SR (27°20'N and 112°16'W) is a mining port intensively impacted by copper exploitation since the late 19th century, with approximately 370 million tons of waste disposed on land and into adjacent coastal waters (Shumilin et al. 2011). This long-term anthropogenic impact has resulted in high concentrations of metals in sediments, such as Cu (3,856 mg kg⁻¹ dry weight), Zn (2,599 mg kg⁻¹), Co (635 mg kg⁻¹), Mn (6,906 mg kg⁻¹), Pb (236 mg kg⁻¹), Cd (240 mg kg⁻¹) and U (11.8 mg kg⁻¹),




Figure 1. Geographic location of the study sites of the mussel *Modiolus capax* in the Gulf of California. La Paz Bay includes Las Tarabillas, San Juan de la Costa, and Punta Prieta.

exceeding established ecotoxicological quality guidelines (Shumilin et al. 2013, Jonathan et al. 2016). In addition, numerous studies have documented elevated concentrations of metals in the tissues of marine species from this site (Gutiérrez-Galindo et al. 1999, Cadena-Cárdenas et al. 2009, Muñoz-Barbosa & Huerta-Díaz 2013, Roldán-Wong et al. 2018, Yee-Duarte et al. 2018, 2024, 2020, Martínez-Ayala et al. 2022, Eraso-Ordoñez 2023, Pintueles-Tamayo et al. 2025), including the target species of the present study, which has been reported to accumulate up to 11 times higher concentrations of various elements compared to organism from less impacted areas (Gutiérrez-Galindo et al. 1999). Consistently, geoaccumulation index assessments categorize this area as strongly and extremely polluted (classes 4 to 6) for elements such as Mn, Ba, Cd, Co, Mo, Pb, Sr, and Zn, reflecting the long-term influence of historical mining and smelting activities in the area (Jonathan et al. 2016). The local climate is arid, with a mean annual temperature and precipitation of 23.4°C and 117.2 mm, respectively

(Rodríguez-Figueroa 2004). The mean sea surface temperature (SST) is 22.4°C, with a mean pH of 8.04 (Góngora-Gómez et al. 2022), and chlorophyll-*a* (Chl-*a*) concentrations ranging from 0.84 to 10.21 mg m⁻³ (Yee-Duarte 2017).

Second, we considered three sites in LPB: a) Punta Prieta (PP) (24°22'N and 110°30'W), an area adjacent to a thermoelectric plant and a port for fuel unloading and storage, in which high concentrations of total polycyclic aromatic hydrocarbons (PAHs) have been reported (Roldán-Wong et al. 2020); b) San Juan de la Costa (SJC) (24°39'N and 110°68'W) an area with Cd, Pb, Fe, Ni, Cu, and Zn in sediments derived from phosphorite mining, which is still in operation to date, the mouth of the San Juan river, and natural deposits located northwest of San José Island (Escobedo-Fregoso 2006, Méndez et al. 2006, Rodríguez-Castañeda et al. 2006, Cadena-Cárdenas et al. 2009, SGM 2019); c) Las Tarabillas (TS) (24°64'N and 110°69'W), a site with the presence of Cd and other metals as a result of the upwelling of the seafloor

towards the surface as a natural process due to its proximity to the San Gregorio rock formation (Segovia-Zavala et al. 2003, Méndez et al. 2006, Cantú-Medellín et al. 2009). The LPB climate is desert, with mean annual precipitation below 200 mm (Coen & Cruz-Bello 2019). The mean sea temperature in LPB is 26.5°C, with a mean pH of 7.95 (Góngora-Gómez et al. 2022), while Chl-*a* concentrations range from 0.8 to 2.1 mg m⁻³ (Coria-Monter et al. 2017).

Specimen collection

At each site, mussel specimens were collected by free diving at depths of 2-5 m between 8:00 and 10:00 h during February, June, September, and November in the period 2017-2020. Subsequently, we transported the specimens in seawater to the laboratory for processing, following bioethical and regulatory guidelines for marine animals used in scientific research (Crook & Walters 2011, Crespi-Abril & Rubilar 2021).

Upon arrival at the laboratory, the biometric data of each mussel was recorded, including shell length (SL, maximum distance along the anterior-posterior axis). (± 0.1 mm), total weight (TW), wet weight (WW, weight without shell), and shell weight (SW) (± 0.1 g).

Size frequency, condition index, and weight-length relationship

The size distribution of *M. capax* at each site was analyzed by constructing frequency histograms from SL. The nutritional status, as a physiological condition of a specimen, was evaluated using the CI with the following equation (Mouneyrac et al. 2008):

$$CI = (WW / TW) \times 100$$

To improve CI estimation and comparison among sites, only organisms in the size range of 6.3 to 9.6 cm SL were used; this range was represented at each site.

As an additional indicator of the health status of *M. capax*, the type of growth of mussels was evaluated by relating the weight (TW, WW, SW) of each specimen to its length, through a WLR analysis using the potential model $P = \alpha L^b$ (Le Cren 1951) and adjusting the parameters for maximum likelihood from the logarithmic transformation:

$$\ln(P) = \ln(\alpha) + b \times \ln(SL)$$

where α is the intercept or initial growth coefficient, b is the slope or allometry coefficient that indicates the type of growth (Gaspar et al. 2001, Froese 2006); P is

TW, WW, or SW (in their respective model), and SL is shell length.

Environmental characterization

Data on SST (°C) and Chl-*a* concentration (mg m⁻³) were obtained through the NASA GIOVANNI application using the MODIS Aqua instrument at a spatial resolution of 4 km. To assess similarity and differences in SST and food availability among study sites, interannual time series spanning 2017-2020 were plotted for each site.

The influence of anthropogenic and natural contamination at each site was characterized using published information documenting long-term environmental conditions in the region, including evidence of historical mining activities, reported trace metal concentrations, and other contaminants in sediments and biota from previous studies. This information was used to provide an environmental context for each site and to support comparisons among sites with contrasting disturbance scenarios. SST, Chl-*a* concentrations, and documented pollution sources were treated as descriptive environmental variables and used as a preliminary framework to interpret spatial differences in mussel organismal condition.

Statistical analysis

Data normality was assessed using the Kolmogorov-Smirnov test with Lilliefors correction and homoscedasticity with Levene's test (Zar 2010). All statistical comparisons among sites, relationships, Chl-*a* concentrations, temperatures, and indices were performed using the Kruskal-Wallis test followed by Dunn's multiple comparison test. To detect and remove outliers, a graph was constructed for each comparison, which improved the estimation of the WLR (Froese 2006, De la Hoz et al. 2016) and the other analyses without altering the data distribution.

Regarding the WLR, a Student's *t*-test was used to determine whether there was a significant difference between the allometry coefficient (b) and the growth hypothesis ($H_0: b = 3, H_1: b \neq 3$) (Zar 2010). Confidence intervals were calculated for α and b with a 95% confidence level using the accelerated bootstrap method with bias correction, and 1,000 resamples (Efron 1987, Zar 2010). Additionally, the degree of association among the variables was determined using the coefficient of determination (R^2) to measure goodness of fit (De la Hoz et al. 2016). All statistical tests were performed in R (R Core Team 2022), at a 95% significance level (Zar 2010).

RESULTS

Environmental variables

Although some fluctuations were observed, the statistical analyses revealed no significant interannual differences in SST or Chl-*a* concentrations within individual sites ($P > 0.05$). Therefore, the mean values calculated throughout the study period were used to represent the environmental conditions at each site in subsequent comparisons. Significant differences in mean SST were detected among sites ($H_{(3)} = 13.1$, $P = 0.004$). SR showed the lowest temperature of all sites ($\bar{x} = 24.3 \pm 4.5^\circ\text{C}$) with ranges of 16 to 31°C , whereas PP ($26.0 \pm 3.2^\circ\text{C}$), SJC ($26.5 \pm 3.2^\circ\text{C}$), and TS ($26.2 \pm 3.3^\circ\text{C}$) were statistically similar ($P > 0.05$), with ranges between 21 and 32°C .

On the other hand, significant differences in mean Chl-*a* concentration were detected among sites ($H_{(3)} = 71.37$, $P < 0.001$). SR exhibited the highest Chl-*a* concentration ($\bar{x} = 1.61 \pm 0.87 \text{ mg m}^{-3}$), followed by PP ($0.99 \pm 0.38 \text{ mg m}^{-3}$, $P < 0.001$). In contrast, SJC and TS showed the lowest concentrations and did not differ significantly from each other (0.78 ± 0.49 , $0.69 \pm 0.41 \text{ mg m}^{-3}$, respectively) ($P > 0.05$) (Fig. 2).

Mussel size

A total of 691 specimens were collected, distributed as follows: 272 from SR, 136 from SJC, 167 from PP, and 116 from TS. Significant differences among sites were recorded in all the *M. capax* biometric parameters evaluated ($H_{(3)} = 241.49$, $P < 0.001$). Significantly larger (SL) and heavier (TW) specimens were collected at SR ($\bar{x} = 9.4 \pm 1.5 \text{ cm SL}$; $\bar{x} = 117.0 \pm 52.8 \text{ g TW}$, respectively), followed by SJC ($7.8 \pm 2.0 \text{ cm}$; $75.9 \pm 48.3 \text{ g}$) and TS ($7.4 \pm 1.2 \text{ cm}$; $64.6 \pm 28.1 \text{ g}$). On the other hand, the mussels collected at PP were the smallest and lightest ($\bar{x} = 6.7 \pm 1.4 \text{ cm}$; $\bar{x} = 46.6 \pm 27.4 \text{ g}$). The heaviest mussels (WW) were recorded in SR ($\bar{x} = 25.9 \pm 10.0 \text{ g WW}$), followed by those collected at SJC ($\bar{x} = 16.7 \pm 9.9 \text{ g}$), while those with the lowest biomass were collected at TS ($12.6 \pm 5.8 \text{ g}$), followed by PP ($10.0 \pm 5.9 \text{ g}$) (Fig. 3).

Size frequency, condition index, and weight-length relationship

The size-frequency distribution of *M. capax* by site is shown in Figure 4. Most mussels with sizes greater than 8.5 cm SL were collected in SR, followed by those collected in SJC (7.4 to 9.6 cm SL), while most specimens collected in PP and TS had sizes between 4.1 and 8.5 cm SL.

The variation in the *M. capax* CI showed significant differences among the study sites ($H_{(3)} = 65.67$, $P < 0.001$). Mussels from SR (CI = $24.1 \pm 5.76\%$) and SJC ($23.2 \pm 4.79\%$) showed the best nutritional status, followed by PP ($21.7 \pm 3.77\%$), while TS specimens had the lowest nutritional status ($18.9 \pm 2.95\%$) (Fig. 5).

The analysis of the WLR of *M. capax* showed isometric growth ($H_0: b = 3$, i.e. a proportional increase in length and weight) in most sites (Student's *t*-test, $P > 0.05$), except in SR and TS, where one of the WLRs (TW-SL, WW-SL, and SW-SL) showed a greater increase in length than in WW (negative allometric growth). PP was the only site where *M. capax* showed negative allometric growth ($b < 3$) in all WLR (Student's *t*-test, $P < 0.05$) (Table 1). The coefficient of determination (R^2) was greater than 0.9, confirming the strong relationship between weight and length.

DISCUSSION

Temperature and food availability are key environmental factors that regulate multiple physiological responses in organisms (Storero et al. 2012). Additionally, anthropogenic disturbance and local habitat quality can influence organismal condition through complex pathways (Cajaraville et al. 2000, Rouabhi et al. 2019). Mussels are widely recognized as effective biological models for environmental studies because they are sessile, widely distributed, long-lived, can accumulate pollutants, and are sensitive to environmental variation (Widdows et al. 2002, Dellali et al. 2023).

The present study recorded marked spatial differences in the biometric parameters (SL, TW, WW) of *M. capax* across the study sites, allowing the identification of three distinct groups. Mussels from SR were consistently the largest and heaviest, followed by SJC and TS mussels, while PP mussels were the smallest and lightest. These spatial differences align with contrasting environmental conditions among sites. SR recorded the highest Chl-*a* concentrations (1.61 mg m^{-3}) and the lowest temperature (24.3°C , with a minimum of 16°C) compared with the other study sites. In bivalves, such as *Mytilus galloprovincialis*, temperatures between 10 and 20°C combined with elevated food availability enhance filtration rates and promote somatic growth (Filgueira et al. 2009, Kholodov et al. 2017, Joyce et al. 2019), allowing organisms to increase visceral mass and improve nutritional condition (Mathieu & Lubet 1993, Lee et al.




Figure 2. Variation in sea surface temperature and chlorophyll-*a* concentrations across study sites of *Modiolus capax* in the Gulf of California during the period 2017-2020. Data distributions are shown with violin plots. Black horizontal lines indicate the mean, and colored circles show data scatter. Different letters denote statistically significant differences among sites based on Dunn's *post-hoc* test ($P < 0.05$).

2015). This pattern matches the highest WW (25.9 g) recorded in SR. However, despite their greater visceral mass, mussels from SR showed a negative allometric relationship between visceral mass and SL, reflected in a more elongated shell morphology and a lower proportional visceral mass. Shell shape influences biometric relationships and may affect CI estimates, as discussed below.

In contrast, sites in LPB (SJC, TS, and PP) had higher temperatures, lower Chl-*a* concentrations, and smaller, lighter mussels. Notably, despite broadly similar temperatures and food availability conditions across LPB sites, significant differences in biometric parameters were still detected among them. Mussels from SJC and TS had comparable shell sizes and TWs; however, SJC specimens showed higher biomass than those from TS and even PP, a site with slightly higher food availability than SJC and TS. This difference may relate to the high phosphorus levels in SJC's coastal waters, attributed to its proximity to the phosphorite mine (Cantú-Medellín et al. 2009, Chávez-Sánchez et al. 2017), which produces up to 15,000 t (SGM 2019). Although nutrients were not directly measured in this

study, the higher visceral mass and isometric growth recorded in SJC mussels suggest a distinct nutrient enrichment pattern relative to TS, where negative allometry between visceral mass and SL was recorded. In this sense, similar relationships among phosphorus availability, visceral mass, and reproductive investment have been reported for other mussel species, including *Lampsilis radiata siliquoidea* (Nalepa et al. 1991) and *M. demissus* (Kuenzler 1961), indicating that nutrient enrichment may influence visceral mass condition and growth patterns in bivalves.

However, although PP had the second-highest Chl-*a* concentration, only surpassed by SR, mussels from this site were the smallest and lightest (lowest visceral mass weight) of all sites analyzed. This apparent decoupling between food availability and biometric performance suggests that factors other than trophic conditions may be constraining overall size at this site. One potential factor influencing mussel size and visceral yield at PP is local population density. Field observations during sampling indicated high mussel densities in a relatively small area, a condition known to intensify intraspecific competition for food and space,




Figure 3. Biometric parameters of *Modiolus capax* across study sites in the Gulf of California. Boxplots and violin charts, along with colored circles representing individual data points for each locality, display data distributions. Within each box, the horizontal line represents the median and the central circle indicates the mean. Horizontal bars at the top denote statistically significant differences among sites based on Dunn's post-hoc test ($P < 0.05$).

generate overcrowding, and negatively affect somatic growth in bivalves (Mallet & Carver 1991, Alunno-Bruscia et al. 2001, Chelyadina 2018). This scenario is consistent with the lower biomass observed for a given SL and with the negative allometric relationships recorded across all weight-length parameters at PP. Other mussel populations have shown similar density conditions (Lauzon-Guay et al. 2005).

In addition to biological interactions, local thermal conditions may also contribute to the reduced biometric performance at PP. This site is near a thermoelectric power plant that discharges heated cooling water into a coastal environment, a process documented in other coastal regions (Martínez-Córdoba 1985, Cardoso-Mohedano et al. 2015). Such thermal discharges can produce abrupt fluctuations in SST, potentially inducing acute thermal stress in resident organisms (Masanja et al. 2023). In this regard, elevated temperatures and thermal variability reduce somatic growth (Hiebenthal et al. 2012, Rahman & Rahman 2021) and induce sublethal or lethal effects in bivalves (Masanja et al. 2023, Smith et al. 2023). Evidence from other regions of the GC supports the sensitivity of *Modiolus* species to extreme thermal conditions. For instance, thermal discharges from the Puerto Libertad thermoelectric plant have been associated with localized seawater temperature rises above 34°C and with massive mortalities of *Modiolus* spp. At the same time, other benthic invertebrates (clams and cirripedia) persisted under similar conditions (Martínez-Córdoba 1985). These observations suggest that mussels may be more vulnerable to acute thermal variations relative to other invertebrate taxa.

Chemical stressors may also contribute to the environmental context at PP. Previous studies have reported elevated concentrations of polycyclic aromatic hydrocarbons (PAHs) in mussels from this site, with levels (173.39 ng g⁻¹ dw) exceeding those measured in surrounding sediments (Roldán-Wong et al. 2020) and comparable to values reported for moderately impacted coastal environments (see Table 3 in Baumard et al. 1998). In this context, the co-occurrence of thermal alterations, documented chemical exposure, and high population density suggests that mussels at PP are exposed to a complex set of environmental pressures. Collectively, these factors provide a plausible framework for interpreting the reduced biometric performance observed at PP, without attributing the observed patterns to any single stressor.




Figure 4. Size frequency distributions of *Modiolus capax* across study sites in the Gulf of California.

Chronic metal exposure is another important environmental baseline across these sites, as at least three study areas have documented elevated metal concentrations (Méndez et al. 2006, Shumilin et al. 2013). At SR, tissue copper levels in *M. capax* have been reported to exceed other regions of the Baja California peninsula baselines up to 11-fold (Gutiérrez-Galindo et al. 1999); nevertheless, SR mussels maintained the largest shell sizes and visceral masses in this study, indicating that chronic metal presence does not necessarily result in stunted somatic growth. Other mussel species in polluted environments have shown similar patterns, where somatic growth and size structure did not decline in parallel with contaminant loads (Chelomin et al. 1998). Such responses are consistent with the physiological and anatomical features that contribute to regulation and detoxification of trace elements, including highly specialized organs involved in metal binding and clearance, such as the kidney and the byssal gland system (through the byssus) (Yonge 1962, Lobel 1986, Szefer et al. 2002, Yap et al. 2003, Chelyadina et al. 2022). The relevance of these traits becomes evident when contrasting *M. capax* with other bivalves co-occurring at SR, such as *Megapitaria squalida* (Yee-Duarte et al. 2017) and *Dosinia ponderosa* (Eraso-Ordoñez 2023), which

possess a much smaller kidney (*pers. obs.*) and lack the byssal gland (Yonge 1962), exhibiting deteriorated condition and stunted somatic growth under similar environmental conditions.

The above suggests that the reduced biometric parameters of mussels collected from LPB sites are more related to the high temperatures and low food availability than to previously reported metal concentrations in SJC (Escobedo-Fregoso 2006, Méndez et al. 2006, SGM 2021) and near TS (between 3 and 8 km) (Segovia-Zavala et al. 2003, Méndez et al. 2006, Cantú-Medellín et al. 2009). However, as noted above, findings of PP mussels with high total PAH concentrations (Roldán-Wong et al. 2020) highlight the need for additional studies to investigate the combined effects of these chemicals and thermal stress at this site.

Morphological plasticity in shell shape has been documented in bivalves exposed to contrasting environmental conditions, including sites with elevated metal concentrations, where variation in shell proportions or convexity has been reported (Chelyadina & Popov 2018, Chelyadina et al. 2022, Sepúlveda et al. 2024). A remarkable example was reported for *Macoma balthica*, in which greater elongation and convexity were observed in organisms that bioaccumu-




Figure 5. Spatial variation in the condition index of *Modiolus capax* across study sites in the Gulf of California. Data distributions are shown using raincloud plots that combine individual observations (dotplots), boxplots, and kernel density plots for each locality. Different letters denote statistically significant differences among sites based on Dunn's *post-hoc* test ($P < 0.05$).

Table 1. Type of growth of *Modiolus capax* from each study site in the Gulf of California. TW: total weight, SL: shell length, WW: wet weight, SW: shell weight. Bold numbers indicate allometric growth ($H_1: b \neq 3$, Student's *t*-test).

Locality	Relationship	Confidence Interval			Confidence Interval			R^2	Growth type	P -value
		a	Lower	Upper	b	Lower	Upper			
Santa Rosalía	TW-SL	0.28	0.21	0.38	2.63	2.50	2.76	0.94	Isometry	0.097
	WW-SL	0.09	0.07	0.12	2.45	2.34	2.57	0.93	Allometry (-)	0.003
	SW-SL	0.18	0.13	0.26	2.71	2.56	2.87	0.91	Isometry	0.178
Punta Prieta	TW-SL	0.32	0.27	0.38	2.54	2.45	2.63	0.97	Allometry (-)	0.031
	WW-SL	0.06	0.05	0.08	2.59	2.45	2.72	0.95	Allometry (-)	0.008
	SW-SL	0.23	0.18	0.29	2.58	2.47	2.70	0.96	Allometry (-)	0.024
San Juan de la Costa	TW-SL	0.25	0.21	0.30	2.68	2.59	2.77	0.98	Isometry	0.137
	WW-SL	0.05	0.04	0.07	2.66	2.53	2.80	0.97	Isometry	0.145
	SW-SL	0.18	0.15	0.22	2.71	2.61	2.81	0.98	Isometry	0.189
Las Tarabillas	TW-SL	0.26	0.17	0.39	2.70	2.50	2.89	0.93	Isometry	0.098
	WW-SL	0.11	0.07	0.18	2.28	2.04	2.51	0.90	Allometry (-)	0.011
	SW-SL	0.17	0.11	0.26	2.78	2.58	2.99	0.91	Isometry	0.216

lated high concentrations of metals in their tissues, such as Cu ($\sim 100 \text{ mg kg}^{-1}$), Zn ($\sim 690 \text{ mg kg}^{-1}$), Ag ($\sim 3 \text{ mg kg}^{-1}$), and As ($\sim 11 \text{ mg kg}^{-1}$) (Sokolowski et al. 2002). In the present study, mussels from SR were not only the largest and heaviest overall but also showed a negative allometric relationship between biomass and SL, indicating that soft-tissue accumulation did not keep pace with SL, a pattern consistent with shell elongation

and geometric shape changes previously associated with metal exposure. This decoupling between visceral mass and shell elongation affects CI estimation because conventional biometric indices assume isometric or proportional growth between internal cavity volume and visceral mass; therefore, shell shape modifications, such as elongated shell morphology, can artificially yield lower CI values in SR mussels despite their large

visceral mass. Consequently, these findings highlight the need to account for shell morphological variation and length-weight allometric relationships when using biometric indices as proxies for organismal condition in metal-impacted environments. The results indicate that *M. capax* maintains high biometric performance despite chronic exposure to metal concentrations associated with historical mining activity that exceed ecotoxicological guidelines at SR (Shumilin et al. 2013, Jonathan et al. 2016). Paradoxically, mussels from this site were consistently the largest and heaviest among all sites analyzed, contrasting with acute experimental observations. For instance, although Strömberg (1982) documented reduced SL and high mortality in *M. edulis* exposed to high Cu and Zn concentrations, those results were derived from small mussels (19-29 mm SL) under short-term exposure (10-22 days), conditions that differ markedly from chronic, lifelong exposure experienced by adult mussels in wild environments.

This discrepancy suggests that large, heavy *M. capax* individuals persist at SR due to population-level acclimation or phenotypic plasticity, traits widely documented in intertidal bivalves (Bayne 2004, Sousa et al. 2007) under fluctuating and anthropogenically altered environments (Sampaio et al. 2022, Góngora-Gómez et al. 2024). These observations highlight the need to distinguish acute experimental responses from chronic, field-based patterns when assessing organismal sensitivity at polluted sites. Future studies integrating physiological, biochemical, and molecular biomarkers will be necessary to identify tolerance thresholds, detoxification pathways, and potential trade-offs that define the balance between resilience and vulnerability in *M. capax* under multi-stressor coastal environments.

CONCLUSION

This study shows that spatial variability in biometric parameters and organismal condition in *M. capax* across GC coastal environments is driven primarily by the interplay between environmental factors (SST and food availability) and localized anthropogenic pressures. Our findings indicate that shell morphological alterations (such as elongation) induced under potentially stressful anthropogenic conditions decouple visceral mass accumulation from SL. This decoupling suggests a critical methodological limitation, showing that conventional length-based condition indices may artificially underestimate organismal condition unless morphological shape alterations are accounted for.

Consequently, WLR, allometric patterns, and morphometrically adjusted condition indices serve as robust, non-invasive ecological baselines for evaluating bivalve population status across coastal environments. Finally, to establish direct cause-effect mechanisms under complex multi-stressor scenarios, future research should integrate these population-level biometric baselines with physiological and biochemical biomarkers in anthropogenically impacted marine ecosystems.

ACKNOWLEDGMENTS

F.A. Aguilar-Santana is a recipient of scholarships from BEIFI (No. A230184), SECIHTI (No. 784901) and the Fundación TELMEX-Telcel (No. 232039532); the results presented herein are part of his PhD Thesis. M. Arellano-Martínez received grants from COFAA, EDI, and SNII- SECIHTI. We thank María Elena Sánchez-Salazar for editing the English manuscript.

Funding

This study was funded by the Secretaría de Investigación y Posgrado of the Instituto Politécnico Nacional through SIP grants 20241674, 20250240, and 20260064.

Credit author contribution

F.A. Aguilar-Santana: conceptualization, data curation, formal analysis, investigation, methodology, validation, writing-original draft, writing-review and editing, A.R. Rivera-Camacho: methodology, fieldwork, I.S. Racotta: validation, writing-review and editing, M. Arellano-Martínez: conceptualization, formal analysis, funding acquisition, methodology, project administration, resources, supervision, validation, writing-original draft, writing-review and editing.

Conflict of interest

The authors declare no conflicts of interest.

REFERENCES

- Alunno-Bruscia, M., Bourget, E. & Fréchette, M. 2001. Shell allometry and length-mass-density relationship for *Mytilus edulis* in an experimental food-regulated situation. *Marine Ecology Progress Series*, 219: 177-188.
- Álvarez-Borrego, S. 2010. Physical, chemical and biological oceanography of the Gulf of California. In: Brusca, R.C. (Ed.). *The Gulf of California: biodiversity and conservation*. The University of Arizona Press, Tucson, pp. 22-48.

- Anestis, A., Pörtner, H.O., Lazou, A., et al. 2008. Metabolic and molecular stress responses of sublittoral bearded horse mussel *Modiolus barbatus* to warming sea water: implications for vertical zonation. *Journal of Experimental Biology*, 211: 2889-2898.
- Assan, D., Kuebutornye, F.K.A., Mustapha, U.F., et al. 2020. Effects of climate change on marine organisms. *American Journal of Climate Change*, 9: 204-216. doi: 10.4236/ajcc.2020.93013
- Baumard, P., Budzinski, H. & Garrigues, P. 1998. Polycyclic aromatic hydrocarbons in sediments and mussels of the western Mediterranean Sea. *Environmental Toxicology and Chemistry*, 17: 765-776. doi: 10.1002/etc.5620170501
- Bayne, B.L. 2004. Phenotypic flexibility and physiological trade-offs in the feeding and growth of marine bivalve molluscs. *Integrative and Comparative Biology*, 44: 425-432. doi: 10.1093/icb/44.6.425
- Brander, K., Botsford, L., Ciannelli, L., et al. 2010. Human impacts on marine ecosystems. In: Barange, M., Field, J.G., Harris, R.P., et al. (Eds.). *Marine ecosystems and global change*. Oxford Academic, Oxford, pp. 41-72. doi: 10.1093/acprof:oso/9780199558025.003.0003
- Brusca, R.C. 2010. Introduction. In: Brusca, R.C. (Ed.). *The Gulf of California: Biodiversity and conservation*. University of Arizona Press, Arizona, pp. 1-6. doi: 10.2307/j.ctt181hwrc.4
- Brusca, R.C. & Hendrickx, M.E. 2010. Invertebrate biodiversity and conservation in the Gulf of California. In: Brusca, R.C. (Ed.). *The Gulf of California: Biodiversity and conservation*. University of Arizona Press, Arizona, pp. 72-95. doi: 10.2307/j.ctt181hwrc.8
- Cadena-Cárdenas, L., Méndez-Rodríguez, L., Zenteno-Savín, T., et al. 2009. Heavy metal levels in marine mollusks from areas with, or without, mining activities along the Gulf of California, Mexico. *Archives of Environmental Contamination and Toxicology*, 57: 96-102. doi: 10.1007/s00244-008-9236-0
- Cajaraville, M.P., Bebianno, M.J., Blasco, J., et al. 2000. The use of biomarkers to assess the impact of pollution in coastal environments of the Iberian Peninsula: a practical approach. *Science of the Total Environment*, 247: 295-311. doi: 10.1016/S0048-9697(99)00499-4
- Canosa, L.F. & Bertucci, J.I. 2023. The effect of environmental stressors on growth in fish and its endocrine control. *Frontiers in Endocrinology*, 14: 1109461. doi: 10.3389/fendo.2023.1109461
- Cantú-Medellín, N., Olguín-Monroy, N.O., Méndez-Rodríguez, L.C., et al. 2009. Antioxidant enzymes and heavy metal levels in tissues of the black chocolate clam *Megapitaria squalida* in Bahía de La Paz, Mexico. *Archives of Environmental Contamination and Toxicology*, 56: 60-66. doi: 10.1007/s00244-008-9156-z
- Cardoso-Mohedano, J.G., Bernardello, R., Sanchez-Cabeza, J.A., et al. 2015. Thermal impact from a thermoelectric power plant on a tropical coastal lagoon. *Water, Air & Soil Pollution*, 226: 1-11. doi: 10.1007/s11270-014-2202-8
- Chávez-Sánchez, T., Piñón-Gimate, A., Serviere-Zaragoza, E., et al. 2017. Recruitment in *Ulva* blooms in relation to temperature, salinity and nutrients in a subtropical bay of the Gulf of California. *Botanica Marina*, 60: 257-270. doi: 10.1515/bot-2016-0066
- Chelomin, V.P., Belcheva, N.N. & Zakhartsev, M.V. 1998. Biochemical mechanisms of adaptation to cadmium and copper ions in the mussel *Mytilus trossulus*. *Russian Journal of Marine Biology*, 24: 330-336.
- Chelyadina, N.S. 2018. Phenotypic and sexual structure of *Mytilus galloprovincialis* Lam., cultivated on the mussel-oyster farm in the outer harbor of Sevastopol city (Crimea, Black Sea). *Marine Biological Journal*, 3: 86-93. doi: 10.21072/mbj. 2018.03.3.09
- Chelyadina, N.S. & Popov, M.A. 2018. Comparative analysis of the allometric characteristics of the mussel *Mytilus galloprovincialis* from different habitats (the Crimea, the Black Sea). *Bulletin of Voronezh State University Series: Chemistry. Biology. Pharmacy*, 2: 264-269.
- Chelyadina, N.S., Popov, M.A., Pospelova, N.V., et al. 2022. Effects of heavy metals on sex inversion of the mussel *Mytilus galloprovincialis* Lam., 1819 in the coastal zone of the Black Sea. *Marine Pollution Bulletin*, 185: 114323. doi: 10.1016/j.marpolbul.2022. 114323
- Clark, M.S., Peck, L.S. & Thyrring, J. 2021. Resilience in Greenland intertidal *Mytilus*: the hidden stress defense. *Science of the Total Environment*, 767: 144366. doi: 10.1016/j.scitotenv.2020.144366
- Coen, M.A. & Cruz-Bello, G.M. 2019. Living with risk: Climate change and vulnerability. *Community perceptions in peri-urban areas of La Paz City, Mexico*. *Sociedad y Ambiente*, 19: 109-136. doi: 10.31840/sya.v0i19.1936
- Coria-Monter, E., Monreal-Gómez, M.A., De León, D.A.S., et al. 2017. Wind-driven nutrient and subsurface chlorophyll-*a* enhancement in the Bay of La Paz, Gulf of California. *Estuarine, Coastal and*

- Shelf Science, 196: 290-300. doi: 10.1016/j.ecss.2017.07.010
- Crespi-Abril, A.C. & Rubilar, T. 2021. Moving forward in the ethical consideration of invertebrates in experimentation: Beyond the three R's principle. *Revista de Biología Tropical*, 69: 346-357. doi: 10.15517/rbt.v69isuppl.1.46366
- Crook, R.J. & Walters, E.T. 2011. Nociceptive behavior and physiology of molluscs: animal welfare implications. *ILAR Journal*, 52: 185-195. doi: 10.1016/j.neuropharm.2022.109370
- De La Hoz-M, J., Motta, J. & Paramo, J. 2016. Length-weight relationships for 36 deep-sea fish in the Colombian Caribbean Sea. *Journal of Applied Ichthyology*, 32: 1356-1359. doi: 10.1111/jai.13217
- Dellali, M., Mardassi, K., Harrath, A.H., et al. 2023. Physiological responses of the bivalves *Mytilus galloprovincialis* and *Ruditapes decussatus* following exposure to Phenanthrene: toxicokinetics, dynamics and biomarkers study. *Animals*, 13: 151. doi: 10.3390/ani13010151
- Doney, S.C., Ruckelshaus, M., Emmett-Duffy, J., et al. 2012. Climate change impacts on marine ecosystems. *Annual Review of Marine Science*, 4: 11-37. doi: 10.1146/annurev-marine-041911-111611
- Efron, B. 1987. Better bootstrap confidence intervals (with discussion). *Journal of the American Statistical Association*, 82: 171-200.
- Eraso-Ordoñez, J.F. 2023. Análisis de la condición fisiológica, histopatológica y reproductiva de *Dosinia ponderosa* (Mollusca: Bivalvia) en el Puerto Minero de Santa Rosalía, BCS, México. Ph.D. Thesis, CICIMAR-IPN, La Paz.
- Escobedo-Fregoso, C. 2006. Evaluación de la síntesis de metalotioneínas inducidas por cadmio en la almeja *Megapitaria squalida* de Bahía de La Paz B.C.S. Tesis de Licenciatura, Universidad Autónoma de Baja California Sur, La Paz.
- Filgueira, R., Fernández-Reiriz, M.J. & Labarta, U. 2009. Clearance rate of the mussel *Mytilus galloprovincialis*. I. Response to extreme chlorophyll ranges. *Ciencias Marinas*, 35: 405-417.
- Filgueira, R., Comeau, L.A., Landry, T., et al. 2013. Bivalve condition index as an indicator of aquaculture intensity: a meta-analysis. *Ecological Indicators*, 25: 215-229. doi: 10.1016/j.ecolind.2012.10.001
- Froese, R. 2006. Cube law, condition factor and weight-length relationships: History, meta-analysis and recommendations. *Journal of Applied Ichthyology*, 22: 241-253. doi: 10.1111/j.1439-0426.2006.00805.x
- García-Corona, J.L., Rodríguez-Jaramillo, C., Saucedo, P.E., et al. 2018. Internal energy management associated with seasonal gonad development and oocyte quality in the horse mussel *Modiolus capax* (Bivalvia; Mytilidae). *Journal of Shellfish Research*, 37: 475-483. doi: 10.2983/035.037.0302
- García-Domínguez, F.A., Carvalho-Saucedo, L. & Vélez-Arellano, N. 2018. Reproductive cycle of two *Modiolus capax* (Bivalvia: Mytilidae) beds at different depths at Ensenada de La Paz, Gulf of California, Mexico. *Hidrobiológica*, 28: 313-323. doi: 10.24275/uam/izt/dcb/hidro/2018v28n3/Garcia
- Gaspar, M., Santos, V. & Vasconcelos, P. 2001. Weight-length relationships of 25 bivalve species (Mollusca: Bivalvia) from the Algarve coast (southern Portugal). *Journal of the Marine Biological Association of the United Kingdom*, 81: 805-807.
- Góngora-Gómez, A.M., Arellano-Martínez, M., Rodríguez-Figueroa, G.M., et al. 2022. Variaciones morfológicas de la concha del mejillón choro, *Modiolus capax*, en dos poblaciones silvestres del suroeste del Golfo de California. *Revista de Biología Marina y Oceanografía*, 57: 148-161. doi: 10.22370/rbmo.2022.57.3.4069
- Góngora-Gómez, A., Hernández-Sepúlveda, J., Isola, T., et al. 2024. Phenotypic plasticity of shell shape and growth of mussels to environmental conditions: *Mytella strigata* in the southeast Gulf of California. *Latin American Journal of Aquatic Research*, 52: 690-702. doi: 10.3856/vol52-issue5-fulltext-3128
- Gosselin, L.A. & Chia, F.S. 1995. Characterizing temperate rocky shores from the perspective of an early juvenile snail: the main threats to survival of newly hatched *Nucella emarginata*. *Marine Biology*, 122: 625-635.
- Gutiérrez-Galindo, E.A., Villaescusa-Celaya, J.A. & Arreola-Chimal, A. 1999. Bioaccumulation of metals in mussels from four sites of the coastal region of Baja California. *Ciencias Marinas*, 25: 557-578. doi: 10.7773/cm.v25i4.726
- Helmholz, H., Ruhnau, C., Profrock, D., et al. 2016. Seasonal and annual variations in physiological and biochemical responses from transplanted marine bioindicator species *Mytilus* spp. during a long-term field exposure experiment. *Science of the Total Environment*, 565: 626-636. doi: 10.1016/j.scitotenv.2016.04.202
- Hiebenthal, C., Philipp, E., Eisenhauer, A., et al. 2012. Interactive effects of temperature and salinity on shell formation and general condition in Baltic Sea *Mytilus*

- edulis* and *Arctica islandica*. Aquatic Biology, 14: 289-298. doi: 10.3354/ab00405
- Horwitz, R., Norin, T., Watson, S.A. et al. 2020. Near-future ocean warming and acidification alter foraging behaviour, locomotion, and metabolic rate in a keystone marine mollusk. Scientific Reports, 10: 5461. doi: 10.1038/s41598-020-62304-4
- Johan, O., Bengen, D.G., Zamani, N.P., et al. 2015. The distribution and abundance of black band disease and white syndrome in Kepulauan Seribu, Indonesia. HAYATI Journal of Biosciences, 22: 105-112. doi: 10.1016/j.hjb.2015.09.001
- Jonathan, M.P., Shumilin, E., Rodríguez-Figueroa, G.M., et al. 2016. Potential toxicity of chemical elements in beach sediments near Santa Rosalía copper mine, Baja California Peninsula, Mexico. Estuarine, Coastal and Shelf Science, 180: 91-96. doi: 10.1016/j.ecss.2016.06.015
- Joyce, P.W.S., Kregting, L.T. & Dick, J.T.A. 2019. Relative impacts of the invasive Pacific oyster, *Crassostrea gigas*, over the native blue mussel, *Mytilus edulis*, are mediated by flow velocity and food concentration. NeoBiota, 45: 19-37.
- Kholodov, V.I., Pirkova, A.V. & Ladygina, L.V. 2017. Cultivation of mussels and oysters in the Black Sea. LLC Izdat-Print, Voronezh.
- Kuenzler, E.J. 1961. Structure and energy flow of a mussel population in a Georgia salt marsh. Limnology and Oceanography, 6: 191-204.
- Lauzon-Guay, J.S., Hamilton, D.J. & Barbeau, M.A. 2005. Effect of mussel density and size on the morphology of blue mussels (*Mytilus edulis*) grown in suspended culture in Prince Edward Island, Canada. Aquaculture, 249: 265-274. doi: 10.1016/j.aquaculture.2005.03.048
- Le Cren, E.D. 1951. The length-weight relationship and seasonal cycle in gonad weight and condition in the perch (*Perca fluviatilis*). Journal of Animal Ecology, 20: 201-219. doi: 10.2307/1540
- Lee, Y.J., Choi, K.S., Lee, D.S., et al. 2015. The role of the adductor muscle as an energy storage organ in the pen shell *Atrina japonica* (Reeve, 1858). Journal of Molluscan Studies, 81: 502-511. doi: 10.1093/mollus/eyv025
- Lobel, P.B. 1986. Role of the kidney in determining the whole soft tissue zinc concentration of individual mussels (*Mytilus edulis*). Marine Biology, 92: 355-359. doi: 10.1007/bf00392676
- López-Carvallo, J.A., Saucedo, P.E., Rodríguez-Jaramillo, C., et al. 2017. Carbohydrate-rich diets meet energy demands of gametogenesis in hatchery-conditioned mussels (*Modiolus capax*) at increasing temperatures. Journal of Shellfish Research, 36: 649-657. doi: 10.2983/035.036.0314
- Lucas, A. & Beninger, P.G. 1985. The use of physiological condition indices in marine bivalve aquaculture. Aquaculture, 44: 187-200. doi: 10.1016/0044-8486(85)90243-1
- Mallet, A.L. & Carver, C.E. 1991. An assessment of strategies for growing mussels in suspended culture. Journal of Shellfish Research, 10: 471-477.
- Martínez-Ayala, J.C., Galván-Magaña, F., Tripp-Valdez, A., et al. 2022. Heavy metal concentrations in the Pacific sharpnose shark *Rhizoprionodon longurio* from the Santa Rosalia mining zone, Baja California Sur, Mexico. Marine Pollution Bulletin, 182: 114018. doi: 10.1016/j.marpolbul.2022.114018
- Martínez-Córdova, L.R. 1985. Estudio ecológico previo al establecimiento de una planta termoeléctrica en Puerto Libertad, Sonora, México. Ciencias Marinas, 11: 93-111. doi: 10.7773/cm.v11i1.451
- Masanja, F., Yang, K., Xu, Y., et al. 2023. Impacts of marine heat extremes on bivalves. Frontiers in Marine Science, 10: 1159261. doi: 10.3389/fmars.2023.1159261
- Mathieu, M. & Lubet, P. 1993. Storage tissue metabolism and reproduction in marine bivalves a brief review. Invertebrate Reproduction & Development, 23: 123-129. doi: 10.1080/07924259.1993.9672303
- Méndez, L., Palacios, E., Acosta, E., et al. 2006. Heavy metals in the clam *Megapitaria squalida* collected from wild and phosphorite mine impacted sites in Baja California, Mexico: considerations for human health effects. Biological Trace Element Research, 110: 275-287. doi: 10.1385/bter:110:3:275
- Mercado-Silva, N. 2005. Condition index of the eastern oyster, *Crassostrea virginica* (Gmelin, 1791) in Sapelo Island, Georgia: effects of site, position on bed and pea crab parasitism. Journal of Shellfish Research, 24: 121-126. doi: 10.2983/0730-8000 (2005)24[121:CIOTEO]2.0.CO;2
- Moullec, F., Asselot, R., Auch, D., et al. 2021. Identifying and addressing the anthropogenic drivers of global change in the North Sea: a systematic map protocol. Environmental Evidence, 10: 1-10. doi: 10.1186/s13750-021-00234-y
- Mouneyrac, C., Linot, S., Amiard, J., et al. 2008. Biological indices, energy reserves, steroid hormones and sexual maturity in the infaunal bivalve *Scrobicularia plana* from three sites differing by their level of contamination. General and Comparative

- Endocrinology, 157: 133-141. doi: 10.1016/j.ygcen.2008.04.010
- Muñoz-Barbosa, A. & Huerta-Díaz, M.A. 2013. Trace metal enrichments in nearshore sediments and accumulation in mussels (*Modiolus capax*) along the eastern coast of Baja California, Mexico: Environmental status in 1995. *Marine Pollution Bulletin*, 77: 71-81. doi: 10.1016/j.marpolbul.2013.10.030
- Nalepa, T., Gardner, W. & Malczyk, J. 1991. Phosphorus cycling by mussels (Unionidae: Bivalvia) in Lake St. Clair. *Hydrobiologia*, 219: 239-250. doi: 10.1007/BF00024758
- Páez-Osuna, F., Álvarez-Borrego, S., Ruiz-Fernández, A.C., et al. 2017. Environmental status of the Gulf of California: a pollution review. *Earth-Science Reviews*, 166: 181-205. doi: 10.1016/j.earscirev.2017.01.014
- Pintueles-Tamayo, J.F., Marmolejo-Rodríguez, A.J., Galván-Magaña, F., et al. 2025. Bioaccumulation and biomagnification of arsenic in the muscle and liver of the speckled guitarfish *Pseudobatos glaucostigmus* in Santa Rosalía, Gulf of California, Mexico. *Marine Pollution Bulletin*, 211: 117356. doi: 10.1016/j.marpolbul.2024.117356
- Przeslawski, R., Davis, A.R. & Benkendorff, K. 2005. Synergistic effects associated with climate change and the development of rocky shore molluscs. *Global Change Biology*, 11: 515-522. doi: 10.1111/j.1365-2486.2005.00918.x
- R Core Team. 2022. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna.
- Rahman, M.S. & Rahman, M.S. 2021. Effects of elevated temperature on prooxidant-antioxidant homeostasis and redox status in the American oyster: Signaling pathways of cellular apoptosis during heat stress. *Environmental Research*, 196: 110428. doi: 10.1016/j.envres.2020.110428
- Ramesha, M.M. & Sophia, S. 2015. Morphometry, length-weight relationships and condition index of *Parreysia favidens* (Benson, 1862) (Bivalvia: Unionidae) from Seeta River in the Western Ghats, India. *Indian Journal of Fisheries*, 62: 18-24.
- Rodríguez-Castañeda, A.P., Sánchez-Rodríguez, I., Shumilin, E.N., et al. 2006. Element concentrations in some species of seaweeds from La Paz Bay and La Paz Lagoon, south-western Baja California, Mexico. *Journal of Applied Phycology*, 18: 399-408. doi: 10.1007/s10811-006-9040-z
- Rodríguez-Figueroa, G.M. 2004. Geoquímica de los oligoelementos, elementos mayores y elementos de las tierras raras, en los sedimentos marinos del distrito minero de Santa Rosalía, B.C.S., México. Tesis de Maestría, CICIMAR-IPN, La Paz.
- Roldán-Wong, N.T., Kidd, K.A., Ceballos-Vázquez, B.P., et al. 2020. Polycyclic aromatic hydrocarbons (PAHs) in mussels (*Modiolus capax*) from sites with increasing anthropogenic impact in La Paz Bay, Gulf of California. *Regional Studies in Marine Science*, 33: 100948. doi: 10.1016/j.rsma.2019.100948
- Rouabhi, Y.L., Grosjean, P., Boutiba, Z., et al. 2019. Reproductive cycle and follicle cleaning process of *Mytilus galloprovincialis* (Mollusca: Bivalvia) from a polluted coastal site in Algeria. *Invertebrate Reproduction & Development*, 63: 255-267. doi: 10.1080/07924259.2019.1631221
- Sampaio, L., Moreira, J., Rubal, M., et al. 2022. A review of coastal anthropogenic impacts on mytilid mussel beds: effects on mussels and their associated assemblages. *Diversity*, 14: 409. doi: 10.3390/d14050409
- Segovia-Zavala, J.A., Delgadillo-Hinojosa, F., Vidal-Talamantes, R., et al. 2003. *Mytilus californianus* transplanted as upwelling bioindicators to two areas off Baja California, Mexico. *Ciencias Marinas*, 29: 665-675.
- Sepúlveda, C.H., Sotelo-Gonzalez, M.I., Osuna-Martínez, C.C., et al. 2024. Shell shape, allometry, and heavy metals content of two oyster species in the southeastern Gulf of California. *Hidrobiológica*, 34: 95-105. doi: 10.24275/jfsu3315
- Servicio Geológico Mexicano (SGM). 2019. Panorama minero del estado de Baja California. SGM, Hidalgo.
- Servicio Geológico Mexicano (SGM). 2021. Panorama minero del estado de Baja California. SGM, Hidalgo.
- Shumilin, E., Gordeev, V., Rodríguez-Figueroa, G., et al. 2011. Assessment of geochemical mobility of metals in surface sediments of the Santa Rosalía mining region, western Gulf of California. *Archives of Environmental Contamination and Toxicology*, 60: 8-25. doi: 10.1007/s00244-010-9532-3
- Shumilin, E., Jiménez-Illescas, Á.R. & López-López, S. 2013. Anthropogenic contamination of metals in sediments of the Santa Rosalía Harbor, Baja California Peninsula. *Bulletin of Environmental Contamination and Toxicology*, 90: 333-337. doi: 10.1007/s00128-012-0923-1
- Smith, K.E., Burrows, M.T., Hobday, A.J., et al. 2023. Biological impacts of marine heatwaves. *Annual Review of Marine Science*, 15: 119-145. doi: 10.1146/annurev-marine-032122-121437

- Sokolowski, A., Fichet, D., Garcia-Meunier, P., et al. 2002. The relationship between metal concentrations and phenotypes in the Baltic clam *Macoma balthica* (L.) from the Gulf of Gdansk, southern Baltic. *Chemosphere*, 47: 475-484. doi: 10.1016/S0045-6535(02)00002-4
- Sousa, R., Freire, R., Rufino, M., et al. 2007. Genetic and shell morphological variability of the invasive bivalve *Corbicula fluminea* (Müller, 1774) in two Portuguese estuaries. *Estuarine, Coastal and Shelf Science*, 74: 166-174.
- Storero, L.P., Narvarte, M.A. & González, R.A. 2012. Reproductive traits of the small Patagonian octopus *Octopus tehuelchus*. *Helgoland Marine Research*, 66: 651-659. doi: 10.1007/s10152-012-0298-z
- Strömberg, T. 1982. Effect of heavy metals (Zn, Hg, Cu, Cd, Pb, Ni) on the length growth of *Mytilus edulis*. *Marine Biology*, 72: 69-72. doi: 10.1007/BF00393949
- Szefer, P., Frelek, K., Szefer, K., et al. 2002. Distribution and relationships of trace metals in soft tissue, byssus and shells of *Mytilus edulis trossulus* from the southern Baltic. *Environmental Pollution*, 120: 423-444. doi: 10.1016/S0269-7491(02)00111-2
- Widdows, J., Donkin, P., Staff, F., et al. 2002. Measurement of stress effects (scope for growth) and contaminant levels in mussels (*Mytilus edulis*) collected from the Irish Sea. *Marine Environmental Research*, 53: 327-356. doi: 10.1016/S0141-1136(01)00120-9
- Yap, C.K., Ismail, A., Tan, S.G., et al. 2003. Can the byssus of greenlipped mussel *Perna viridis* (Linnaeus) from the west coast of Peninsular Malaysia be a biomonitoring organ for Cd, Pb and Zn? Field and laboratory studies. *Environment International*, 29: 521-528. doi: 10.1016/S0160-4120(03)00008-4
- Yee-Duarte, J.A. 2017. Salud Reproductiva de la almeja chocolata *Megapitaria squalida* (Sowerby, 1835) (Mollusca: Bivalvia) en el Puerto Minero de Santa Rosalía BCS, México. Tesis de Maestría, CICIMAR-IPN, La Paz.
- Yee-Duarte, J.A., Arellano-Martínez, M., Roldán-Wong, N.T., et al. 2024. Reduction in reproductive activity from degeneration of testicular follicles in *Megapitaria squalida* (Mollusca: Bivalvia) exposed to metal pollution in the Gulf of California. *Marine Pollution Bulletin*, 205: 116648. doi: 10.1016/j.marpolbul.2024.116648
- Yee-Duarte, J.A., Ceballos-Vázquez, B.P., Arellano-Martínez, M., et al. 2018. Histopathological alterations in the gonad of *Megapitaria squalida* (Mollusca: Bivalvia) inhabiting a heavy metal-polluted environment. *Journal of Aquatic Animal Health*, 30: 144-154. doi: 10.1002/aah.10015
- Yee-Duarte, J.A., Racotta, I.S., Camacho-Mondragón, M.A., et al. 2020. Contrasting reproductive health of female clams *Megapitaria squalida* from two nearby metal-polluted sites in the Gulf of California: Potential effects of copper, lead, and cobalt. *Marine Pollution Bulletin*, 160: 111583. doi: 10.1016/j.marpolbul.2020.111583
- Yonge, C.M. 1962. On the primitive significance of the byssus in the Bivalvia and its effects on evolution. *Journal of the Marine Biological Association of the United Kingdom*, 42: 113-125. doi: 10.1017/S0025315400004495
- Zar, J.H. 2010. Biostatistical analysis. Pearson Prentice Hall, Upper Saddle River.

Received: March 11, 2026; Accepted: September 24, 2026